

2004 GALVESTON BAY INVASIVE SPECIES RISK ASSESSMENT

INVASIVE SPECIES SUMMARY

Created by: Environmental Institute of Houston, University of Houston-Clear Lake
and the Houston Advanced Research Center

Common Name: Red imported fire ant (RIFA)
Latin Name: <i>Solenopsis invicta</i>
Category: Terrestrial Animal
Place of Origin: Brazil
Place of Introduction: "The RIFA was first introduced from Brazil into either Mobile, AL or Pensacola, FL between 1933 and 1945. The black imported fire ant is confined to northeastern Mississippi and northwestern Alabama. However, the RIFA encompasses eleven southeastern states from North Carolina and Texas to Florida, and has recently been found in New Mexico, Arizona, and southern California (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm)."
Date of Introduction: Between 1933 and 1945 (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm)
<p>Life History: "The lifespan of RIFA workers depends on their size. Minor workers may live 30 to 60 days, media workers 60 to 90 days, major workers 90 to 180 days, and queens may live two to six years. Complete lifecycle from egg to adult takes between 22 and 38 days (Hedges 1997).</p> <p>Mating flights are the primary means of colony propagation, secondarily, budding can occur in which a portion of a colony becomes an autonomous unit. After the colony reaches one year of age, reproductive alates are produced. Six to eight mating flights consisting of up to 4,500 alates each occur between the spring and fall (Vinson & Sorenson 1986). Mating flights usually occur midday on a warm (>74°F/24°C), sunny day following rain (Hedges 1997). Mating occurs during flight and the males die soon after mating with females. In the southern United States, as many as 97,000 queens may be produced per acre of infested land per year (Vinson & Sorenson 1986). Alates are often attracted to swimming pools where homeowners can find thousands of winged ants trapped on the water's surface (Hedges 1997).</p> <p>After mating flights occur, it is common to find newly-mated queens clustered together under shelter. This clustering and the cooperation of the newly-mated queens aid in establishing a colony. However, as the colony grows all but one queen will be killed, except in the instance of multiple queen colonies (Vinson & Sorenson 1986).</p> <p>Once the female has mated, she will rake her legs forward to snap her wings off at the basal suture and find a suitable spot to begin a new colony (Holldobler 1990). Often this spot is under rocks, leaves or in a small crack or crevice, such as at the edge of a sidewalk, driveway, or street. The queen will burrow into the soil to excavate a small chamber, which is sealed off to keep predators out. Within 24 hours of mating, the queen will have laid 10 to 15 eggs, which will hatch in eight to 10 days. By the time the first group of eggs hatch, the queen will have laid from 75 to 125 more eggs. The larval stage typically lasts six to 12 days and the pupal stage for nine to 16 days. The newly-mated queen will stop laying eggs until the first batch of workers mature. This process takes from two weeks to one month. The queen will feed the first batch of young larvae oils regurgitated from her crop, trophic eggs or secretions from her salivary glands. The queen's wing muscles, which are no longer needed, break down to provide the nutrients for the young larvae (Vinson & Sorenson 1986).</p> <p>The first group of workers to emerge is characteristically small due to the limitations in nutrients that the queen provides. These workers, termed "minims," burrow out of the chamber and begin foraging for food to feed the queen and new larvae. The minims also begin construction of the mound. Within one month, larger workers are being produced and the mound is growing in size. By six months the colony has reached several thousand workers and the mound can be seen in a field or lawn. Colonies of this size generally contain a few large workers (major workers), many medium sized workers (media workers), and a majority of small workers (minor workers). The three types of workers are all sterile females and serve to perform tasks necessary to maintain the colony. The queen is the single producer of eggs and is capable of producing as many as 1,500 eggs per day. Mature RIFA colonies may contain as many as 240,000 workers with a typical colony consisting of 80,000 workers (Vinson & Sorenson 1986).</p> <p>The diet of foraging workers consists of dead animals, including insects, earthworms, and vertebrates. Workers also collect honeydew and will forage for sweets, proteins, and fats in homes. They are sometimes attracted to piles of dirty laundry. Larvae are fed only a liquid diet until they reach the third instar. When the larvae reach the fourth instar, they are able to digest solid foods. Worker ants will bring solid food rich in protein and deposit it in a depression in front of the mouth of the larvae. The larvae will secrete digestive enzymes that break down the solid food and regurgitate it back to worker ants. The queen is fed some of the digested protein to support egg production. As long as food is plentiful, egg production is at its maximum (Vinson & Sorenson 1986).</p> <p>The sting of the RIFA possesses venom of an alkaloid nature, which exhibits potent necrotoxic activity. Approximately 95% of the venom is composed of these alkaloids, which are responsible for both the pain and white pustule that appears approximately one day after the sting occurs. The remainder of the venom contains an aqueous solution of proteins, peptides, and other small molecules that</p>

produce the allergic reaction in hypersensitive individuals. Worker fire ants will attach to the skin using their mandibles and will subsequently lower the tip of the gaster to inject the stinger into the victim. Thus, fire ants both bite and sting, but only the sting is responsible for the pain and pustule (Goddard 1996).

Multiple Queen Colonies

Early studies of imported fire ant biology indicated that colonies contained single queens (monogyne). However, since 1973, reports have become more frequent of the occurrence of multiple queen (polygyne) colonies. Multiple queen colonies have been found in areas of Florida, Mississippi, Louisiana, Texas, and Georgia, being more frequent in the western edge of their range (Vinson & Sorenson 1986).

Multiple queen colonies differ from single queen colonies in several ways: (1) the mounds are closer together and more numerous per acre of land, (2) the colonies have smaller workers with fewer major workers present, (3) workers are not aggressive towards neighboring colonies, and (4) the queens weigh less and produce fewer eggs than single queens. The overall number of eggs produced in a multiple queen colony is higher than in a single queen colony due to the presence of many queens producing eggs simultaneously. Additionally, fewer reproductive alates of fire ants are produced with the males often being sterile and the females weighing less (Hedges 1998, Vinson & Sorenson 1986) (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm)."

Growth/Size: Workers are between 2.4 and 6 mm long.

Feeding Habits/Diet: "The diet of foraging workers consists of dead animals, including insects, earthworms, and vertebrates. Workers also collect honeydew and will forage for sweets, proteins, and fats in homes. They are sometimes attracted to piles of dirty laundry. Larvae are fed only a liquid diet until they reach the third instar. When the larvae reach the fourth instar, they are able to digest solid foods. Worker ants will bring solid food rich in protein and deposit it in a depression in front of the mouth of the larvae. The larvae will secrete digestive enzymes that break down the solid food and regurgitate it back to worker ants. The queen is fed some of the digested protein to support egg production. As long as food is plentiful, egg production is at its maximum (Vinson & Sorenson 1986).

In agriculture, the RIFA frequently invades soybean crops and heavy infestations invariably yield fewer soybeans (Lofgren & Adams 1981). . . Opportunistic feeding on the young tender growth by red imported fire ant occurs throughout the year and can cause significant damage not only to soybean crops, but citrus, corn, okra, bean, cabbage, cucumber, eggplant, potato, sweet potato, peanut, sorghum, and sunflower (Stimac & Alves 1994). Monetary losses occur when RIFA interfere with the root system of the plant, mechanically disable combine operation, or feed on the young growth of plants to reduce stands (Stimac & Alves 1994). (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm)"

Habitat: "The red imported fire ant builds mounds in almost any type of soil, but prefers open, sunny areas such as pastures, parks, lawns, meadows and cultivated fields. Mounds can reach 18 inches in height, depending on the type of soil. Often mounds are located in rotting logs and around stumps and trees. Colonies also can occur in or under buildings.

Colonies frequently migrate from one site to another. The queen needs only half a dozen workers to start a new colony. They can develop a new mound several hundred feet away from their previous location almost overnight. Flooding causes colonies to leave their mounds and float until they can reach land to establish a new mound. Colonies also can migrate to indoor locations (<http://fireant.tamu.edu/antfacts/biology.html>)."

Attitude (aggressive, etc.): "When a mound is disturbed, ants emerge aggressively to bite and sting the intruder. A white pustule usually appears the next day at the site of the sting (Cohen 1992)."

"Red imported fire ants have been reported to reduce ground-nesting populations of rodents and birds. In certain instances, the RIFA may completely eliminate ground-nesting species from a given area (Vinson & Sorenson 1986). Because there is a 10 to 20 year lapse before reductions in bird populations are observed, it has been suggested that actual effects of the RIFA on animal populations may be underestimated (Mount 1981).

The human toll from RIFA stings is an important public health concern. Stings may produce a large range of reactions from localized pain and swelling to anaphylactic shock, making it hard to estimate the cost to public health (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm)."

Physical Description: "The pedicel, or "waist" in the RIFA consists of two segments. Workers consist of many sizes (polymorphic) between 2.4 to 6 mm (1/8 to 1/4 in) (Hedges 1998). The mandible has four distinct teeth and the antennae are 10-segmented, ending in a two-segmented club. A sting is present at the tip of the gaster. Body color is usually red to brown in color with a black gaster (Hedges 1997).

Mounds are built of soil and are seldom larger than 46 cm (18 in) in diameter. When a mound is disturbed, ants emerge aggressively to bite and sting the intruder. A white pustule usually appears the next day at the site of the sting (Cohen 1992).

It is important to distinguish between the RIFA and the native fire ant in order for appropriate control measures to be taken. Mounds of *S. geminata* will contain workers with square-shaped heads that are larger in proportion to the rest of their body. These workers collect and mill seeds for the colony (Drees 1997). Workers of *S. invicta* do not have workers with disproportionate head to body

ratios (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm).”

Management Recommendations / Control Strategies: include references for existing site-specific strategies

“Fire ants frequently invade home lawns, school yards, athletic fields, golf courses, parks and other recreational areas. Additionally, electrical equipment and utility housing, home gardens, compost piles, mulched flowerbeds, pavement cracks, and the perimeter of bodies of water must all be considered when choosing a method of control.

Two approaches can be taken to effectively manage imported fire ants. Single mound treatments or area-wide broadcast applications usually manage red imported fire ant populations.

Individual Mound Treatments

There are many methods of treating individual mounds. The main advantage to this process is that there are so many choices available to homeowners and pest control operators, though few may actually eliminate the colony. However, the largest disadvantage is that individual mounds must each be located in order to be treated. Individual mound treatments are most beneficial when there are native ants in the same area as imported fire ants. Re-infestation of any treated area, whether by broadcast treatment or individual mound treatment may occur. Six different methods of individual mound treatment are available (Vinson & Sorenson 1986).

1. Mound Drenches. Large volumes of liquid toxic to ants are poured over a mound. Liquids can range from using several gallons of hot water to insecticides mixed with several gallons of water. This method may not reach the queen, who may be deep in the nest, thus preventing colony elimination.
2. Surface Dusts. This method is very similar to mound drenches. A dust or granular insecticide is applied over the top of the mound and then watered into the soil.
3. Mound Injections. The use of insecticides that may be pressurized and injected into a mound. Often this method is more expensive, but more effective, than mound drenches. However, more time may be required for this method and leakage of the insecticide by the equipment may be hazardous to the handler. Again, the queen may not be affected and thus re-infestation may occur.
4. Baits. Baits can be used for both individual mound and broadcast applications. A small amount of the bait is sprinkled around the mound and the ants then forage and bring the bait back to the colony to feed on. This method is slower acting, but more effective than drenching, dusting, or fumigating a mound because the workers will feed the bait to the queen and brood, thus gaining effective control of the colony.
5. Mechanical Control. Certain mechanical and electrical devices are on the market for controlling fire ants, but the efficacy has not been documented (Vinson & Sorenson 1986).
6. Home Remedies. Many homeowners will choose to pour boiling water or ignite flammable liquids over a mound. While these methods may bring about control, they are not recommended because they are both very dangerous, not only to humans, but also to the environment.

Broadcast Treatments

Presently, there are only a few products available for broadcast treatment of large areas. These products are either granular insecticides or baits composed of soybean oil and toxicant on a corn grit carrier. These granules are broadcast over a large area and, upon discovery by the ants, are carried to the colony and fed to nest mates and the queen. This is a very effective treatment but does present problems because (1) some of the bait may be dropped where the ants may not find it, (2) some colonies are well fed so do not feed upon the bait, (3) some baits are light sensitive (as with hydramethylnon) and may inactivate before discovery by the ants, and (4) the baits are not specific to the imported fire ant (Vinson & Sorenson 1986).

Biological Control

Current research is leading the way for introduction of natural enemies of the red imported fire ant. Two hopeful pathogens are the microsporidian protozoan *Thelohania solenopsae* and the fungus *Beauveria bassiana*. Two parasitoid flies from South America, *Pseudacteon tricuspidis* and *Pseudacteon curvatus* have been introduced into the southern states. These flies decapitate worker ants in the final stages of larval development. In addition, a parasitic ant, *Solenopsis daguerri* (Santschi) which invades fire ant colonies and replaces the queen to take control of the colony, is being investigated for another possible biological control agent of the red imported fire ant (http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm).”

References (includes journals, agency/university reports, and internet links):

1. http://creatures.ifas.ufl.edu/urban/ants/red_imported_fire_ant.htm).
2. <http://www.ceris.purdue.edu/napis/pests/ifa/index.html#maps>
3. <http://www.ars.usda.gov/is/AR/archive/sep99/ant0999.htm>. USDA ARS News and Information. Ouch! The Fire Ant Saga Continues.
4. <http://fireant.tamu.edu/> and several pages within this site, including <http://fireant.tamu.edu/antfacts/biology.html>. Texas Imported Fire Ant Research and Management Plan. Texas A&M University, Department of Entomology, College Station.
5. Hedges, S. A. 1997. Handbook of Pest Control, 8th Ed. (D. Moreland, ed.) pp. 531-535. Mallis Handbook and Technical Training

Company.

6. Vinson, S. B.; A. A. Sorenson. 1986. Imported Fire Ants: Life History and Impact. The Texas Department of Agriculture. P. O. Box 12847, Austin, Texas 78711.
7. Holldobler, B.; E. O. Wilson. 1990. The Ants. pp. 165. Belknap Press.
8. Goddard, J. 1996. Physicians Guide To Arthropods of Medical Importance. pp. 374. CRC Press, NY.
9. Lofgren, C. S.; C. T. Adams. 1981. Reduced yield of soybeans in fields infested with the red imported fire ant, *Solenopsis invicta* Buren. [Florida Entomol.](#) 64: 199-202.
10. Stimac, J. L.; S. B. Alves. 1994. Pest Management in the Subtropics: Biological Control A Florida Perspective. (D. Rosen, F. D. Bennett, J. L. Capinera, ed) pp. 353-380. Intercept Limited, Andover, Hants SP10 1 YG, UK.
11. Cohen, P. R. 1992. Imported fire ant stings: clinical manifestations and treatment. *Pediatr. Determol.* 9: 44-48.
12. Mount, R. H. 1981. The red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae) as a possible serious predator on some southeastern vertebrates: direct observations and subjective impressions. *J. Alabama Acad. Sci.* 52: 71-78.
13. Hedges, S. A. 1998. Field Guide for the Management of Structure Infesting Ants, 2nd Ed.(D. Moreland, ed.) pp. 202-216. G. I. E. Publishers, Cleveland, Ohio
14. Drees, B. M. 1997. We're all on the same team when tracking fire ants. <http://fireant.tamu.edu/> (22 June 2000).

Available Mapping Information:

1. Cooperative Agriculture Pest Survey Program, Purdue. <http://www.ceris.purdue.edu/napis/pests/ifa/index.html#maps>
2. Texas Imported Fire Ant Research and Management Plan. Texas A&M University, Department of Entomology, College Station. <http://fireant.tamu.edu/maps/index.html>

Notes: Texas Imported Fire Ant Research and Management Plan. Texas A&M University, Department of Entomology, College Station (<http://fireant.tamu.edu/plan/>). This website summarizes the State-wide management plan as described below.

Overall Goal

The plan's overall goal is to develop products and procedures to reduce fire ant populations to a level that eliminates this insect as a serious pest-~~in~~ in terms of economic losses and health threats.

A Coordinated Effort

This plan differs from past plans in that this is a coordinated effort among all Texas organizations addressing the fire ant problem through research, educational, and regulatory programs: Texas Agricultural Experiment Station, Texas Agricultural Extension Service, Texas Department of Agriculture, Texas Park and Wildlife Department, Texas Tech University, University of Texas, University of Texas at El Paso, University of Texas at San Antonio and industry leaders.